

FSME Policy and Procedures 2-5
REVISION 0
{DATE}

FSME PROCEDURE FOR INTERACTING WITH THE ADVISORY COMMITTEE ON THE MEDICAL USES OF ISOTOPES DURING DEVELOPMENT OF MAJOR MEDICAL ISSUES

1. PURPOSE:

This Federal and State Materials and Environmental Management Programs (FSME) Policy and Procedure (P&P) defines and documents FSME staff guidance and procedures for interfacing with the Advisory Committee on the Medical Uses of Isotopes (ACMUI) during the development of major medical policy issues including medical rulemakings that will be reviewed by the Commission. ACMUI provides advice, as requested by the Director, Division of Materials Safety and State Agreements (MSSA) on policy and technical issues that arise in regulating the medical use of byproduct material for diagnosis and therapy. The ACMUI also raises issues of concern to the medical community to the U.S. Nuclear Regulatory Commission (NRC).

2. BACKGROUND:

In a Staff Requirements Memo (SRM) dated July 21, 2010 (M100708B-Briefing on Proposed Rule on Part 35 Medical Events Definitions-Permanent Implant Brachytherapy, July 8, 2010), the Commission directed the staff to develop internal guidance that requires the staff to include ACMUI recommendations and dissenting views along with the staff's assessment of the ACMUI recommendations and dissenting views, for all major medical policy issues that are submitted to the Commission, including proposed and final rules.

Major medical use policy issues include changes to the Code of Federal Regulations (i.e., rulemaking), as well as certain changes in medical use licensing and inspection guidance and other medical use policy issues that will require Commission review and/or approval. Determination of the status of a medical issue as a major medical policy issue will be made by the MSSA Director or the Director, Division of Intergovernmental Liaison and Rulemaking (DILR). Major medical policy issues may be legally sensitive or significantly impact the public, patients or human research subjects, medical use licensees or stakeholders, or the Agreement States.

Changing the Code of Federal Regulations (i.e., rulemaking) is a significant action. Adding, deleting, or revising regulations impacts licensees, certificate holders and other stakeholders. For this reason, MSSA staff should draft a regulatory basis as a robust foundation for the rulemaking before a rulemaking begins. The regulatory basis should contain the justification for a rule and describe the technical, legal, or policy information that supports the rulemaking and provides a basis for informed decisions during the rulemaking.

Rules that could affect medical activities would require the MSSA to seek input from the ACMUI prior to initiation of the rulemaking. The level of input may range from very little ACMUI involvement (e.g., a recommendation from the ACMUI to support MSSA's proposed revisions to the regulations) to very extensive ACMUI involvement (e.g., a report with information to be used in the regulatory basis).

If during the rulemaking process, MSSA determines it is necessary to make fundamental changes to the regulatory basis due to issues identified by MSSA or the ACMUI, such changes must be reflected in an amended regulatory basis. This will require that work must stop on the rulemaking and that the amended regulatory basis be provided to the rulemaking working group (WG). If this happens, DILR will inform the Executive Director of Operations (EDO) and the Office of the Secretary of the Commission (SECY) that rulemaking schedule needs to be rebaselined. A formal request to the EDO and SECY for rebaselining will follow when DILR has accepted the amended regulatory basis.

3. ROLES AND RESPONSIBILITIES

3.1 Rulemaking

3.1.1 In DILR, Rulemaking Branches (RB) A, and B, have the overall responsibility for the preparation of rulemaking packages. The rule package includes the *Federal Register* notice (FRN) for the proposed or final rule, as well as the appropriate supporting documents (e.g., a regulatory analysis, an environmental assessment, a backfit analysis, Congressional letters, State liaison letters, Agreement State letters, and a press release). In addition, the Office of Management and Budget (OMB) supporting statements are needed for rules with information collection requirements.

3.1.2 The MSSA, Radioactive Materials Safety Branch (RMSB), has the primary responsibility of seeking input from the ACMUI on any proposed or final rulemakings that may affect medical uses of radioactive materials.

3.2 Other Major Medical Policy Issues

The MSSA, RMSB, has the primary responsibility of seeking input from the ACMUI on any proposal that constitutes a major policy issue that may affect medical uses of radioactive materials and will eventually need review or approval from the Commission.

4. PROCEDURES

4.1 Proposed Rules:

After acceptance of the regulatory basis for a proposed rulemaking, the Branch Chief (RBA or RBB) will assign a project manager (PM) to lead the rulemaking in accordance with FSME P&P 6-10 entitled, "FSME Procedures for Preparation and Review of Rulemaking Packages."

The RB PM will work with the WG members to prepare a draft package for the proposed rule.

A draft version of the proposed rule FRN may be provided electronically to the ACMUI PM at the point at which the rulemaking PM and the MSSA WG member believe it to be substantially complete. The ACMUI PM will distribute the draft proposed rule FRN to the ACMUI members for their review and comment. The ACMUI PM may indicate that the documents are predecisional and cannot be publicly released, unless authorized by NRC staff. In accordance with an SRM

dated October 25, 2007 (SECY-07-0134 Evaluation of the Overall Effectiveness of the Rulemaking Process Improvement Implementation Plan), the FSME Director may authorize the release of draft rule text, statements of consideration, and the technical basis for public review and to hold workshops prior to submission of a proposed rule to the Commission.

ACMUI will review the draft rule and consolidate its comments into an ACMUI position. The ACMUI position may include dissenting views of individual ACMUI members. The ACMUI will be given 90 days to complete its review and provide comments. In some cases, the ACMUI may find that additional discussion or information is required to provide quality opinions and request that more time be allotted. The MSSA Director may grant an extension on a case-by-case basis.

The RB PM, in concert with the WG members, will address comments from the ACMUI in the preparation of the final proposed rule package. If comments are substantive or if the draft rule has significant changes, DILR will reissue the package to the appropriate offices for concurrence. The Commission Paper should include a discussion of the staff's interaction with the ACMUI. The Commission paper must specifically include ACMUI comments/recommendations, any dissenting opinions from ACMUI members, plus staff's assessment of ACMUI recommendations and dissenting views and the basis for incorporating or not incorporating ACMUI recommendations into the proposed rule.

After Commission approval, the proposed rule will be published in the *Federal Register* for public comments. The standard public comment period is 75 days.

4.2 Final Rules:

After the public comment period is over, the RB PM will hold periodic meetings with the WG or with specific members to resolve all of the comments received. The RB PM will review the responses to ensure the comments have been appropriately addressed. The RB PM and WG members should identify any controversial issues to management at an early stage.

A draft version of the final rule FRN may be provided electronically to the ACMUI PM at the point at which the RB PM and the MSSA WG member believe it to be substantially complete. The ACMUI PM will distribute the draft final rule FRN to the ACMUI members for their review. The ACMUI PM may indicate that the documents are predecisional and cannot be publicly released, unless authorized by NRC staff. The SRM dated October 25, 2007 (SECY-07-0134) referenced in 4.1 above authorizes the release of the draft proposed rule documents, the FSME Director may authorize the release of draft final rule documents. ACMUI will review the draft rule and consolidate its comments into an ACMUI position. The ACMUI position may include dissenting views of individual ACMUI members. The ACMUI will be given 90 days to complete its review and provide comments. In some cases, the ACMUI may find that additional discussion or information is required to provide quality opinions and request that more time be allotted. The MSSA Director may grant an extension on a case-by-case basis.

The RB PM, in concert with the WG members, will address comments from the ACMUI, and prepare the final rule package (if the comments are substantive or if

the final rule has significant changes from the draft, DILR will reissue the package to the appropriate offices for concurrence). The Commission paper must specifically include ACMUI comments/recommendations, any dissenting opinions from ACMUI members, plus staff's assessment of ACMUI recommendations and dissenting views and the basis for not incorporating ACMUI recommendations into the final rule.

The MSSA WG member may also seek ACMUI's advice and guidance throughout the rulemaking process. The MSSA WG member and the ACMUI PM may share predecisional documents with the ACMUI to get their input. However, the ACMUI PM must indicate that the documents are predecisional and cannot be publicly released or discussed.

The RB PM will take the lead in briefing the ACMUI, if requested by the MSSA management on the status and progress of any medical rulemakings.

After Commission approval, the final rule will be published in the *Federal Register*.

4.3 Other Major Medical Policy Issues:

Other major policy issues that may affect medical uses of radioactive materials may arise as a result of medical use licensing and inspection findings or emerging issues brought to the staff or Commission attention by the medical community. After identification of a major policy issue that may affect medical uses of radioactive materials (other than rulemaking) that the MSSA intends to take to the Commission for review (e.g., SECY paper on a specific issue or significant licensing or inspection guidance revision for medical use licensees), the ACMUI PM will distribute the documents to the ACMUI members for review and comment. The ACMUI will be given at least 60 days to complete its review and provide comments. MSSA may also elect to obtain earlier ACMUI input by including one or more members of the ACMUI on the staff working group developing the policy issue or document. The ACMUI member serving on the WG may be the Chairman, a member most closely associated with the issue, or a member of an ACMUI subcommittee working on the issue or a related issue. It would be understood, however, that these individual members do not represent the ACMUI's collective opinion.

MSSA should provide the ACMUI's comments or recommendations on a proposed or final major medical policy issue, including dissenting views in the document provided to the Commission. MSSA should also describe how the ACMUI views were considered in the development and finalization of the major medical policy issue.

4.4 ACMUI Meetings:

The ACMUI views will be obtained during either regular or special (e.g., teleconference) ACMUI meetings. The meetings will be open to the public unless they may be closed under the provisions of 10 CFR 9.104, "Closed meetings."

5. ACMUI ENDORSEMENT

The draft procedure was presented to the ACMUI for their review and endorsement during the October 20-21, 2010, and December 13, 2010, ACMUI Meetings. The ACMUI members reviewed the draft procedure, provided comments, and unanimously approved the procedure with comments.

6. DOCUMENT HISTORY

| Version | Description of Change | Responsible Division | Date Last Modified |
|---------|-----------------------|----------------------|--------------------|
| 0 | Initial Procedure | DILR/MSSA | |

November 30, 2010

To: Advisory Committee on the Medical Use of Isotopes
From: David Switzer, MS and Walter Roberts, MD
Subject: Written Statement Re: Patient release following iodine-131 therapy

Thank you for the opportunity to provide a written statement with regard to patient release following the administration of therapeutic quantities of iodine-131.

We have provided services for many years for thyroid cancer patients using iodine-131. We have observed that a large number of these patients have been young mothers, who, if released would have had to contend with maintaining separation from their children. A few patients have been incontinent. A few patients have been confined to psychiatric institutions and not compliant with the usual requirements for release had it been feasible. Other instances could be enumerated.

Under good practice there are patients who do indeed qualify for release during their post administration period. At the same time there are many patients whom we may better serve by confinement in minimal care circumstances. Certainly, many studies have been undertaken that confirm the low risk involved for appropriate release of patients and the information generated by these studies should be taken into consideration.

The bottom line is the authorized users, with competent medical physics support when needed, should be able to independently determine when and if it is appropriate to confine a patient and there should be appropriate reimbursement for any and all such instances where confinement is indicated. Guidance provided by the NRC's Guide on radionuclide therapy has been very useful in determining length and need of confinement.

1
2
3
4 **Advisory Committee on the Medical Use of Isotopes (ACMUI)**
5 **Patient Release Subcommittee Report**
6
7
8

9 **Subcommittee Members:** D. Fisher, Ph.D.; D. Gilley, MPA; S. Langhorst, Ph.D. (Chair); S.
10 Mattmuller, MS, R.Ph, BCNP; O. Suleiman, Ph.D.; B. Thomadsen, Ph.D.; J. Welsh, M.D.; P.
11 Zanzonico, Ph.D.
12

13
14 **Charge:** To evaluate patient release/human research subject release issues; to objectively review
15 and analyze data, which may include state regulations and guidance as well as recommendations in
16 international guidance documents; to provide a statement on the issues, including patient release to
17 other than private residences and an annual rather than per-release limit on radiation doses to others
18 from released individuals; and, if appropriate, to provide recommendations for improvements to
19 existing NRC rules and guidance.
20

21
22 **Summary Statements and Recommendations**
23

- 24 **1.** The medical use of radioactive materials provides important diagnostic and therapeutic tools
25 that have well-recognized health benefits^{1,2,3,4}. Use of radionuclides in medicine and patient
26 access to radionuclide medical procedures, with associated public doses at or below typical
27 environmental background levels, should not be burdened by excessive regulatory controls,
28 including controls that may lead some practitioners to avoid their use or to deliver sub-optimal
29 care (such as multiple lower-administered activity treatments) simply to comply with regulatory
30 dose limits. The Subcommittee affirms that radiation doses to other individuals from
31 radioactivity in released patients⁵ can be safely controlled by:
32
- 33 • the current 10 CFR 35.75 patient release criteria⁶,
 - 34 • licensees' use of scientifically developed dose-based release calculation methods, and
 - 35 patient release instructions based on individual patient circumstances, and
 - 36 • patients' and caregivers' understanding of and adherence to the patient release instructions.

¹ NCRP Commentary No. 11, "Dose Limits for Individuals Who Receive Exposure from Radionuclide Therapy Patients", National Council on Radiation Protection and Measurements, February 1995.

² ICRP Publication 94, "Release of Patients after Therapy with Unsealed Radionuclides", International Commission on Radiological Protection, March 2004.

³ NCRP Report No. 155, "Management of Radionuclide Therapy Patients", National Council on Radiation Protection and Measurements, December 2006.

⁴ IAEA Safety Reports Series No. 63, "Release of Patients after Radionuclide Therapy", International Atomic Energy Agency, 2009.

⁵ Use of the term "patient" in this report is intended to also include human research subject.

⁶ NRC Regulation 10 CFR 35.75, "Release of individuals containing unsealed byproduct material or implants containing byproduct material", Nuclear Regulatory Commission.

37
38 Relevant regulations should not be overly prescriptive because the licensee is best qualified to
39 assess the suitability of individual patients for release post-treatment and to provide
40 personalized guidance to patients to assure compliance with the applicable release criteria.
41

- 42 2. Current 10 CFR 35.75 patient release criteria, along with NRC RIS 2003-04⁷, appropriately
43 balance public safety with patient access to medical treatment.
44
- 45 • Based on NRC conclusions documented in the final rulemaking⁸ and lack of further
46 rulemaking changes to these criteria, the current patient release criteria should continue to be
47 considered as per-release dose limits until modified by future rulemaking.
 - 48 • National and international scientific recommendations on patient release are consistent, in
49 principle and practice, with NRC patient release regulations and guidance.
 - 50 • The NRC per-release 5 mSv (500-mrem) dose limit for any individual is consistent with
51 ICRP and IAEA recommendations for caregivers and other members of the patient's
52 household.
 - 53 • For all other members of the general public, NRC requires the licensee to provide written
54 instructions to the patient on ways to keep radiation dose as low as reasonably achievable, or
55 less than 1 mSv (100 mrem). Specifically, these instructions further protect children,
56 pregnant women, and non-caregivers.
- 57
- 58 3. Current NRC guidance on patient release calculations⁹ overestimates caregiver and public doses
59 because the guidance assumes unrealistically conservative assumptions. The Subcommittee
60 recommends that:
61
- 62 • NRC guidance and assumptions should be updated, with assistance from experts, and should
63 include current information on actual radiopharmaceutical biokinetics and calculated or
64 measured patient dose rates.
 - 65 • Updated scientifically-based tools should be developed to assist licensees in determining and
66 documenting compliance with the patient release criteria.
 - 67 • Reasonable assumptions should be employed for calculating realistic doses to people from a
68 released patient.
 - 69 • In addition to private residences, release scenarios should address patient release to other
70 locations (such as hotels, public transport, public events).
- 71
- 72 4. Current NRC instructions for patient release⁹ should be updated, in conjunction with release
73 calculation methods and assumptions, and the NRC should support research efforts to advance
74 understanding and communication of circumstances that impact patient release decisions,
75 instructions and perceptions.
76

⁷ NRC Regulatory Issue Summary 2003-04 "Use of the Effective Dose Equivalent in Place of the Deep Dose Equivalent in Dose Assessments" (February 13, 2003).

⁸ 62 FR 4120: "Criteria for the Release of Individuals Administered Radioactive Material-Final Rule", NRC Docket No. RIN 3150-AE41, January 29, 1997.

⁹ NRC Regulatory Guide 8.39, "Release of Patients Administered Radioactive Materials", Nuclear Regulatory Commission, April 1997.

77 Scientific Evaluation of Patient/Human Research Subject Release Issues

78
79 Experts in radiation protection^{10,11} apply three fundamental principles to the use of radioactive
80 materials:

- 81
- 82 • The Principle of Justification: Any decision that alters the radiation exposure situation
83 should do more good than harm.
- 84 • The Principle of Optimization of Protection: The likelihood of incurring exposure, the
85 number of people exposed, and the magnitude of their individual doses should all be kept as
86 low as reasonably achievable (ALARA), taking into account economic and societal as well
87 as medical factors.
- 88 • The Principle of Application of Dose Limits: The total dose to any individual from
89 regulated sources in planned exposure situations other than medical exposure of patients
90 should not exceed the appropriate limits specified.

91
92 The appropriate use of radioactive materials in medicine is accepted as doing more good than
93 harm. Exposure to the patient is intentional for the direct medical benefit of the patient. Radiation
94 protection experts oppose dose limits for patients because doing so may compromise the
95 effectiveness of the patient's diagnosis or treatment, and thus do more harm than good. Experts
96 emphasize the physician's informed medical justification for a patient's medical procedure while
97 maintaining the patient's radiation dose as low as reasonably achievable, again taking into account
98 economic and societal as well as medical factors.

99 100 Exposure to Other Individuals from Patients Released from Licensee Control

101
102 Patients undergoing therapeutic medical procedures using radioactive materials become a
103 radiation source that may expose other individuals, and therefore warrant appropriate precautions
104 for limiting doses to those individuals. Patients undergoing diagnostic radiopharmaceutical
105 procedures may also expose other individuals to radiation fields. The likely dose to others from
106 nuclear medicine or implant procedures is low, but not necessarily zero^{12,13}. Individuals most
107 likely to be exposed to a released patient are the patient's family members, or other person caring
108 for or comforting the patient (caregiver), who will be in physical proximity of the patient in the
109 initial days following release. Reducing the need for hospital stays also provide patients, their
110 families and caregivers psychological and emotional benefits of having the patient with them and of
111 lowering their health care costs^{13,14}. This also provides societal benefits by reducing the direct
112 economic costs, and commitment, of medical resources required to retain the patient in a hospital,

¹⁰ NCRP Report No. 116, "Limitation of Exposure to Ionizing Radiation", National Council on Radiation Protection and Measurements, March 1993.

¹¹ ICRP Publication 103, "The 2007 Recommendations of the International Commission on Radiation Protection", March 2007.

¹² ICRP Publication 94, "Release of Patients after Therapy with Unsealed Radionuclides", International Commission on Radiological Protection, March 2004.

¹³ NRC NUREG-1492, "Regulatory Analysis on Criteria for the Release of Patients Administered Radioactive Materials, Final Report", by Stewart Schneider and Stephen A. McGuire, Nuclear Regulatory Commission, April 1996.

¹⁴ 62 FR 4120: "Criteria for the Release of Individuals Administered Radioactive Material-Final Rule", NRC Docket No. RIN 3150-AE41, January 29, 1997.

113 and the indirect costs of a patient's or their employer's lost work time¹⁵. Exposures to other
 114 individuals can be effectively managed by the educated patient (or parent or guardian) after release
 115 if that patient follows the instructions provided by the licensee. These instructions help the patient
 116 to maintain doses to levels comparable to or less than variations in natural background radiation
 117 doses. Given the balance of personal and societal benefits gained, and the ability to maintain doses
 118 to others as low as reasonably achievable levels, the NRC concluded in its final rulemaking that the
 119 benefits outweigh the potential of small increased risks associated with the release of patients
 120 administered radioactive materials^{16,17}.

121

122 Scientific Development of Current NRC Patient Release Criteria

123

124 In the early 1990s, the NRC received three petitions for rule making^{18,19,20} concerning the 10
 125 CFR 35.75 patient release criteria, which at that time included an activity-based limit and 10 CFR
 126 20.1301 public dose limits. In response to these petitions, the NRC initiated rulemaking to change
 127 patient release criteria to dose rate-based limits²¹. The NRC evaluated patient release criteria which
 128 appropriately applied the three fundamental principles previously discussed. The NRC considered
 129 three alternatives in its cost-benefit analysis¹⁵ of the controlling criteria for determining when a
 130 patient may be released from the licensee's control:

131

132 **Alternative 1** – 1 mSv (100 mrem) per year dose limit in 10 CFR 20.1301

133

134 **Alternative 2** – less than 1,110 MBq (30 mCi) or less than 0.05 mSv/h (5 mrem/h) at 1
 135 meter per the activity-based, which was the 1996 version of 10 CFR 35.75²²

136

137 **Alternative 3** – 5 mSv (500 mrem) dose limit

138

139 NRC concluded that **Alternative 3** best served the interest of patients and society¹⁶ for the
 140 following reasons:

141

- 142 1. All of the alternatives were compatible with generally accepted radiation protection
 143 principles.
- 144 2. **Alternative 1** was dismissed due to its excessive economic costs and adverse psychological
 145 impact on patients and their families due to the required patient isolation.

¹⁵ NRC NUREG-1492, "Regulatory Analysis on Criteria for the Release of Patients Administered Radioactive Materials, Final Report", by Stewart Schneider and Stephen A. McGuire, Nuclear Regulatory Commission, April 1996.

¹⁶ NRC SECY 96-100: "Final Amendments to 10 CFR Parts 20 and 35 on Criteria for the Release of Individuals Administered Radioactive Material", Nuclear Regulatory Commission, May 8, 1996.

¹⁷ 62 FR 4120: "Criteria for the Release of Individuals Administered Radioactive Material-Final Rule", NRC Docket No. RIN 3150-AE41, January 29, 1997.

¹⁸ 56 FR 26945: "Carol S. Marcus; Filing of Petition for Rulemaking", NRC Docket No. PRM-20-20, June 12, 1991.

¹⁹ 57 FR 8282: "American College of Nuclear Medicine; Receipt of Petition for Rulemaking", NRC Docket No. PRM-35-10, March 9, 1992; and 57 FR 21043: "American College of Nuclear Medicine; Receipt of Amended Petition for Rulemaking", NRC Docket No. PRM-35-10A, May 18, 1992.

²⁰ 59 FR 37950: "American Medical Association; Petition for Rulemaking", NRC Docket No. PRM-35-11, July 26, 1994.

²¹ 59 FR 30724: "Criteria for the Release of Patient Administered Radioactive Material, Proposed Rule", NRC Docket No. RIN 3150-AE41, June 15, 1994.

²² Also referred to as the "30-mCi rule"

- 146 3. **Alternative 3** was preferred over **Alternative 2** because of its more favorable cost-
 147 effectiveness and more positive psychological impact on patients and their families.
 148 4. Basing patient release criteria on the dose to individuals exposed to a patient provided the
 149 consistent, scientific basis of dose for such decisions that treats all radionuclides on a risk-
 150 equivalent basis. The 30-mCi limit (**Alternative 2**), which may have been appropriate for
 151 iodine-131 under some circumstances, was excessive for some patients and clinical
 152 situations using certain other radionuclides (projected doses would be well below the dose
 153 limit), but inadequate for other situations and radionuclides (projected doses exceed the dose
 154 limit).
 155 5. **Alternative 3** allowed physicians flexibility to *not* have to fractionate therapy doses, leading
 156 to improved effectiveness of treatment for the patient while avoiding unnecessary
 157 hospitalization associated with the 30-mCi rule²³.
 158 6. Reduction of medically unwarranted hospital stays provided emotional benefits to patients
 159 and their families. Allowing earlier reunion of families could improve the patient's state of
 160 mind, which in itself improved the outcome of the treatment and led to the delivery of more
 161 effective health care. At the same time, the opportunity to personally care for a seriously ill
 162 family member was comforting to many individuals.
 163

164 Today, the Subcommittee affirms the thorough analysis found in NUREG-1492 and its rational
 165 evaluation of the three alternatives. The NRC's final decision to implement **Alternative 3** as the
 166 patient release criteria found in 10 CFR 35.75 appropriately balanced the three fundamental
 167 radiation protection principles for use of radioactive materials in medicine.
 168

169 Current National and International Recommendations Regarding Released Patients

170
 171 The National Council on Radiation Protection and Measurements (NCRP) recommendations²⁴
 172 specific to release criteria for radionuclide therapy patients in place at the time NRC established the
 173 current 10 CFR 35.75 release criteria were as follows:
 174

| <u>Other Individual</u> | <u>NCRP²⁴ Recommended Dose Limit</u> |
|---|---|
| Public | 1 mSv/y, but 5 mSv/y may be used for infrequent exposures |
| Patient's Family, Adults | 5 mSv/y, 50 mSv/y with special training |
| Patient's Family, Children and Pregnant Women | 1 mSv/y |

175
 176 The NCRP also concluded in this commentary that "a contamination incident that could lead to a
 177 significant intake of radioactive material is very unlikely"²⁵. The most recent NCRP Report on the
 178 subject maintains those same limits²⁶.

²³ In locations where the 30-mCi rule is in effect, some physicians treat thyroid cancer with multiple administrations of 29.9 mCi of I-131 for no reason other than to avoid hospitalization of patients, thereby treating the patient in a protracted, less therapeutically-effective manner, which can compromise the treatment and, ultimately, the well-being of the patient. When physicians choose to treat thyroid cancer with one administration greater than 30 mCi of I-131, patients can be denied treatment, some for many months, until a private hospital bed is available.

²⁴ NCRP Commentary No. 11, "Dose Limits for Individuals Who Receive Exposure from Radionuclide Therapy Patients", National Council on Radiation Protection and Measurements, February 1995.

179
180
181
182
183
184

The International Commission on Radiation Protection (ICRP) recently updated its recommendations on limiting dose to other individuals from the release of patients after therapy with unsealed radionuclides²⁷. The ICRP recommendations incorporate the concept of dose constraint, rather than a dose limit, as follows:

| <u>Other Individual</u> | <u>ICRP²⁷ Recommendations</u> |
|---|---|
| Public | 1 mSv/y (limit) |
| Relatives, Visitors, and Caregivers | A few mSv/episode (constraint) |
| Infants, Young Children, and Casual Visitor | 1 mSv/y (limit) |

185
186
187
188
189

The International Atomic Energy Agency (IAEA) also recently published a safety series report on the release of radionuclide therapy patients²⁸. The IAEA endorsed the ICRP recommendations and further clarified its criteria in a recent position statement²⁹.

190
191
192
193

All three of the above authoritative national and international advisory bodies agreed that the decision to hospitalize or release a patient should be determined on an individual basis and should be based on dose criteria rather than on residual-activity criteria (as with the previous 30-mCi rule).

194
195
196
197
198
199
200
201
202

The physician's decision should also take into account the patient's wishes and medical condition, his or her physical and mental capacity to understand and follow instructions, occupational and public exposures, family considerations (including the presence of children and pregnant women in the household), cost, and environmental factors. These advisory bodies' recommendations incorporated the concept of maintaining the dose to other individuals as low as reasonably achievable, and recognized the need for flexibility in the regulatory authority's practical application of limits and constraints so that patient physical and psychological factors, as well as economic and societal factors, are properly considered.

203
204
205
206

The ICRP noted that determination of the overall costs associated with various methodologies related to release of patients after therapy with unsealed radionuclides had generally not been attempted²⁷. The ICRP stated:

207
208
209
210

“Ideally, ‘costs’ should include psychological and adverse health consequences, as well as monetary costs. Cost-benefit analysis for a specific issue may vary substantially from country to country, but it does provide a tool that may help the optimization process.”

²⁵ NCRP Commentary No. 11, “Dose Limits for Individuals Who Receive Exposure from Radionuclide Therapy Patients”, National Council on Radiation Protection and Measurements, February 1995.

²⁶ NCRP Report 155, “Management of Radionuclide Therapy Patients.” National Council on Radiation Protection and Measurements, December 2006.

²⁷ ICRP Publication 94, “Release of Patients after Therapy with Unsealed Radionuclides”, International Commission on Radiological Protection, March 2004.

²⁸ IAEA Safety Reports Series No. 63, “Release of Patients after Radionuclide Therapy”, International Atomic Energy Agency, 2009.

²⁹ IAEA Position Statement, “Release of Patients after Radionuclide Therapy”, International Atomic Energy Agency, February 23, 2010.

211 The ICRP cited the NRC’s NUREG-1492 cost-benefit analysis as a scientifically appropriate
 212 example.

213
 214 The Subcommittee finds the current 10 CFR 35.75 release criteria to be consistent with the
 215 practical application of nationally and internationally recommended dose constraints and limits, and
 216 to be in harmony with public safety, humane patient care, and cost-effective delivery of medical
 217 treatment.

218

219 Control of Dose to Other Individuals from Released Patients

220

221 In contrast to diagnostic nuclear medicine procedures, doses to the public, patients’ relatives,
 222 and others may need to be limited after some therapeutic procedures. The preponderance of peer-
 223 reviewed scientific data demonstrate that the radiation dose from internal contamination of other
 224 individuals from released patients is far less significant than that from external exposure^{30,31,32,33}.
 225 Because of its physical properties and the extent of its use, I-131 is the most likely therapeutic
 226 radionuclide having potential to cause radiation dose to medical staff, the public and family
 227 members. Therefore, the Subcommittee has focused its review on circumstances associated with I-
 228 131 therapy patients.

229

230 Prior to patient release, the licensee has responsibilities established by NRC regulations and
 231 license conditions for controlling dose to other individuals exposed to an I-131 therapy patient.
 232 These controls incorporate well-established and straightforward concepts of limiting exposure:
 233 minimizing time, maximizing distance from the source (i.e., the patient), and, to the extent practical,
 234 using shielding. Controls include measures to prevent or at least minimize radioactive
 235 contamination; a medical facility’s use of universal precautions^{34,35} and infection controls^{36,37}
 236 effectively achieve this. The licensee has responsibility to evaluate the circumstances of the
 237 planned patient release to ensure compliance with 10 CFR 35.75³⁸, which permits a licensee to
 238 “authorize the release from its control any individual who has been administered unsealed
 239 byproduct material or implants containing byproduct material if the total effective dose equivalent
 240 to any other individual from exposure to the released individual will not likely exceed 5 mSv (0.5

³⁰ NCRP Commentary No. 11, “Dose Limits for Individuals Who Receive Exposure from Radionuclide Therapy Patients”, National Council on Radiation Protection and Measurements, February 1995.

³¹ NRC NUREG-1492, “Regulatory Analysis on Criteria for the Release of Patients Administered Radioactive Materials, Final Report”, by Stewart Schneider and Stephen A. McGuire, Nuclear Regulatory Commission, April 1996.

³² ICRP Publication 94, “Release of Patients after Therapy with Unsealed Radionuclides”, International Commission on Radiological Protection, March 2004.

³³ IAEA Safety Reports Series No. 63, “Release of Patients after Radionuclide Therapy”, International Atomic Energy Agency, 2009.

³⁴ OSHA Regulation 29 CFR 1910.1030, “Bloodborne Pathogens”, Occupational Safety & Health Administration, Department of Labor.

³⁵ CDC Fact Sheet, “Universal Precautions for Prevention of Transmission of HIV and Other Bloodborne Infections”, Centers for Disease and Control Prevention, Department of Health and Human Services, 1996 update.

³⁶ CDC, “Guidelines for Environmental Infection Control in Health-Care Facilities”, Centers for Disease and Control Prevention, Department of Health and Human Services, 2003.

³⁷ CDC, “2007 Guidelines for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings”, Centers for Disease and Control Prevention, Department of Health and Human Services, 2007.

³⁸ NRC Regulation 10 CFR 35.75, “Release of individuals containing unsealed byproduct material or implants containing byproduct material”, Nuclear Regulatory Commission.

241 rem)”. The licensee is also required to “provide the released individual, or the individual's parent or
242 guardian, with instructions, including written instructions, on actions recommended to maintain
243 doses to other individuals as low as is reasonably achievable if the total effective dose equivalent to
244 any other individual is likely to exceed 1 mSv (0.1 rem)”. This regulatory language characterizes
245 the responsibility of the licensee as ensuring that the dose to an individual from a released patient is
246 not *likely* to exceed the specified dose limit, rather than as certitude that the dose limit will not be
247 exceeded.

248
249 In the case of an orally administered therapeutic radionuclide (such as I-131 sodium iodide),
250 vomiting shortly after its administration is a contamination concern. The NRC concluded in its
251 final rulemaking for the current 10 CFR 35.75³⁹:

252
253 “Vomiting is seldom an important elimination route for radiopharmaceuticals after the patient
254 has left the medical facility since orally administered radiopharmaceuticals such as iodine-131
255 are rapidly absorbed, within a half hour, by the gastrointestinal system.”

256
257 Vomiting is a rare event, and can often be prevented by giving antiemetics to the patient prior to
258 administration of the radionuclide. The risk of vomiting in public can be further mitigated by
259 having the patient remain in a designated monitored area at the facility for a short period of time
260 post-administration, when vomiting is most likely.

261
262 Once an I-131 therapy patient is released, NRC’s regulatory control, and thus the licensee’s
263 responsibilities⁴⁰, ends³⁹. At this point, the patient, parent or guardian assumes responsibility for
264 managing radiation exposure to other individuals based on instructions provided by the licensee.
265 These instructions should be straightforward and easy to follow so that the patient will understand
266 how to minimize radiation doses to other individuals as low as reasonably achievable. Instructions
267 include maintaining distance from other people, minimizing time in public places, measures to
268 reduce the spread of radioactive contamination, and the length of time the patient should follow
269 each such precaution⁴¹. As part of the implementation of the current 10 CFR 35.75 release criteria,
270 the NRC worked with the Society of Nuclear Medicine (SNM) to prepare a pamphlet that provides
271 practical information for patients receiving treatment with radioiodine⁴². The NRC noted in final
272 rulemaking for the current 10 CFR 35.75³⁹ that “American medical practice routinely depends on
273 patients following instructions, such as instructions on when and how to take medications”.

274
275 As a licensee reviews the I-131 therapy patient’s post-release living and traveling
276 circumstances, certain precautions may be emphasized or lengths of time adjusted for special
277 circumstances, such as those involving potential exposure of children or pregnant women or the
278 need to use public transportation to return home or to stay in a hotel or other non-private residence

³⁹ 62 FR 4120: “Criteria for the Release of Individuals Administered Radioactive Material-Final Rule” (NRC Docket No. RIN 3150-AE41), January 29, 1997.

⁴⁰ The term “licensee’s responsibilities” refers only to the control of radioactive material under NRC regulations, and does not include the physician’s continuing responsibilities for medical care of the patient.

⁴¹ NRC Regulatory Guide 8.39, “Release of Patients Administered Radioactive Materials”, Nuclear Regulatory Commission, April 1997.

⁴² SNM Pamphlet, “Guidelines for Patients Receiving Radioiodine Treatment,” Society of Nuclear Medicine, 1997. This pamphlet may be obtained from the Society of Nuclear Medicine, 1850 Samuel Morse Drive, Reston, VA 20190-5316.

279 prior to returning home. As the IAEA noted⁴³, “The success of a patient release program is
280 critically dependent on the quality and specificity of the information provided to the patient, the
281 skill with which it is communicated, and whether or not the patient believes the information
282 provided.” The IAEA also advised that the precautions “should be based upon realistic models of
283 behavior, including realistic occupancy factors, and should not be over-cautious”⁴⁴.

284
285 The NRC adopted a dose-based limit in its final rulemaking because it “better expresses the
286 NRC’s primary concern for the public’s health and safety”⁴⁵. Scientists^{46,47} have measured doses to
287 other individuals, primarily family members and other caregivers, from released I-131 therapy
288 patients, and the actual doses received by these individuals are significantly less than those
289 conservatively projected by the licensee as the basis for the patient release.

291 Use and Misuse of Conservative Assumptions in Estimating Dose to Other Individuals

292
293 With implementation of the current 10 CFR 35.75 release criteria, the NRC issued guidance⁴⁸ to
294 assist licensees with determining when a patient could be released, when instructions to patients
295 were required, and what records must be generated and maintained. NRC guidance on calculating
296 dose to other individuals was primarily based on release of an I-131 therapy patient using what is
297 now judged to be very conservative assumptions^{49,50}. As noted, the IAEA advised that these dose
298 calculations should be realistic and not overly-cautious⁴⁴. Although NRC’s 1997 guidance was
299 conservative, the NRC practice of establishing risk-informed and performance-based regulations⁵¹
300 allowed licensees the practical flexibility to use more reasonable guidance and realistic calculations
301 in determining compliance with the current 10 CFR 35.75 release criteria⁴⁵.

302
303 As previously discussed, licensees must evaluate an I-131 therapy patient’s post-release living
304 circumstances in order to choose reasonable specific calculation assumptions and to provide
305 appropriate instructions specific for that patient. On the other hand, when performing such analyses
306 for a generalized patient population, more conservative assumptions may be chosen to account for a
307 greater range of living or traveling circumstances. And, experts may assume activities, distances,
308 occupancy factors, and so forth, that far exceed values likely to be encountered in practice to

⁴³ IAEA Safety Reports Series No. 63, “Release of Patients after Radionuclide Therapy”, International Atomic Energy Agency, 2009.

⁴⁴ IAEA Position Statement, “Release of Patients after Radionuclide Therapy”, International Atomic Energy Agency, February 23, 2010.

⁴⁵ 62 FR 4120: “Criteria for the Release of Individuals Administered Radioactive Material-Final Rule”, NRC Docket No. RIN 3150-AE41, January 29, 1997.

⁴⁶ Grigsby PW, Siegel BA, Baker S, & Eichling, JO. “Radiation exposure from outpatient radioactive iodine (I-131) therapy for Thyroid Carcinoma”. JAMA. 2000;283:2272–2274.

⁴⁷ Rutar FJ, Augustine SC, Colcher D, et al. “Outpatient treatment with 131I-anti-B1 antibody: radiation exposure to family members”. J Nucl Med. 2001;42:907–915.

⁴⁸ NRC Regulatory Guide 8.39, “Release of Patients Administered Radioactive Materials”, Nuclear Regulatory Commission, April 1997.

⁴⁹ Siegel JA, Marcus CS, Stabin MG, “Licensee Over-Reliance on Conservatism in NRC Guidance Regarding the Release of Patient Treated with I-131”, Health Physics (93:667-677), December 2007.

⁵⁰ ICRP Publication 94, “Release of Patients after Therapy with Unsealed Radionuclides”, International Commission on Radiological Protection, March 2004.

⁵¹ NRC “The Risk-Informed and Performance-Based Plan (RPP)”, <http://www.nrc.gov/about-nrc/regulatory/risk-informed/rpp.html>.

309 thereby demonstrate that if such highly improbable scenarios are compatible with release criteria,
 310 then more realistic dose projections could be expected to be much lower. However, some may
 311 misuse the end result from such extreme calculations uncritically, that is, without consideration of
 312 how unrealistic the underlying assumptions are, and thus precipitate unnecessary public safety
 313 concerns and alarm.

314
 315 An example of such a calculation is found in the latest ICRP recommendations⁵². The ICRP
 316 made this calculation to demonstrate the importance of an I-131 therapy patient taking precautions
 317 to reduce or prevent internal contamination of children and infants. The ICRP's concluding
 318 statements accompanying this calculation are as follows:

319
 320 "Contamination of infants and young children with saliva from a treated patient during the
 321 first few days after radioiodine therapy could result in significant doses to the child's
 322 thyroid, and potentially raise the risk of subsequent radiation-induced thyroid cancer".

323
 324 "Thyroid cancer as a result of contamination (particularly with saliva) may be a significant
 325 risk for those under 20 years of age."

326
 327 As described in Paragraphs (68) and (69) of the ICRP report⁵², the following unrealistic
 328 assumptions were used:

- 329
- 330 • The I-131 therapy patient (parent) does not follow the precautions given in their oral and
 331 written instructions to minimize contact with their own infants and children;
 - 332 • The I-131 therapy patient (parent) transfers 1 milliliter (e.g., approximately ¼ teaspoon) of
 333 saliva (55,500 Bq = 1.5 µCi) by kissing the child in the first day after therapy; and,
 - 334 • The thyroid cancer incidence from this child's calculated thyroid dose is estimated based on
 335 preliminary data of cancer incidence being studied in children who ingested larger amounts
 336 of radioactive iodine and other radionuclides in milk and vegetables contaminated from the
 337 Chernobyl accident⁵³.

338
 339 The ICRP report stated that actual measurements from children when parents followed appropriate
 340 precautions resulted in lower thyroid doses than those indicated by this calculation. In one study⁵⁴,
 341 iodine activity was detected in only 25 of 89 children; even though some of these parents did not
 342 receive, understand, or follow the precautions. So even without proper instruction, 64 of the 89
 343 children had no detectable iodine activity.

344

⁵² ICRP Publication 94, "Release of Patients after Therapy with Unsealed Radionuclides", International Commission on Radiological Protection, March 2004.

⁵³ Another study of children administered diagnostic amounts (5 to 15 µCi) of I-131 found no incidence of cancer – Dickman PW, et. al., "Thyroid Cancer Risk After Thyroid Examination with I-131: a Population-Based Cohort Study in Sweden", *Int. J. Cancer*: 106, 580-587 (2003).

⁵⁴ Barrington, S.F., O'Doherty, M.J., Kettle, A.G., et al. "Radiation Exposure of Families of Outpatients Treated with Radioactive Iodine (iodine-131) for Hyperthyroidism", *Eur. J. Nucl. Med.* 26, 686–692 (1999).

345 The Subcommittee agrees that a released I-131 therapy patient should be instructed to take
346 special precautions to minimize dose to children and pregnant women. The 1997 SNM pamphlet⁵⁵
347 that many licensees provide to their I-131 therapy patients instructs the patient to avoid kissing the
348 first few days following treatment, and to avoid prolonged physical contact, especially with children
349 and pregnant women, explaining that the thyroid glands of children and fetuses are more sensitive
350 to the effects of I-131 than those of adults.

351
352 The NRC issued a Regulatory Issue Summary (RIS)⁵⁶ in 2008, which included the first ICRP
353 concluding statement listed above, but provided no details regarding the assumptions. The RIS also
354 stated:

355
356 “However, as described in the Background section of this RIS, for some I-131 therapies,
357 such as oral administration of sodium iodide I-131, the ICRP cautions that the internal dose
358 to infants and young children who may come in contact with a released patient could be
359 significant.”

360
361 “The guidance recommends that licensees consider not releasing patients, administered I-
362 131, whose living conditions may result in unnecessary exposure of infants and young
363 children.”

364
365 The intent of this RIS was to remind licensees of precautions (established in 1997 with the current
366 10 CFR 35.75 release criteria) that should be discussed with their I-131 therapy patients. The
367 Subcommittee recommends that these types of caution statements should be fully explained, and
368 that future documents of this type should include a statement for patients to consult their physician
369 for additional information specific to their medical procedure.

370 371 Release of I-131 Therapy Patients to Locations other than a Private Residence

372
373 The NRC asked the ACMUI to review a draft RIS being developed to address the release of I-
374 131 therapy patients to locations other than a private residence. As part of the ACMUI’s analysis,
375 the ACMUI Subcommittee calculated the radiation dose to other individuals from release of an I-
376 131 therapy patient to a hotel. Despite the possibility of misunderstanding or misuse of the
377 resulting calculation and conclusions, the Subcommittee used overly conservative assumptions and
378 parameters, along with reasonable ones, to demonstrate that even highly unlikely dose projections
379 do not exceed the release criteria and that reasonable doses are comparable to variations in
380 background radiation doses.

381
382 The example calculations, assumptions used in each case, and the results of this analysis are
383 presented in the Report Appendix⁵⁷. The Subcommittee concluded that when a licensee assesses
384 the I-131 therapy patient’s planned living situation upon release, provides the patient with simple

⁵⁵ SNM Pamphlet, “Guidelines for Patients Receiving Radioiodine Treatment,” Society of Nuclear Medicine, 1997. This pamphlet may be obtained from the Society of Nuclear Medicine, 1850 Samuel Morse Drive, Reston, VA 20190-5316.

⁵⁶ NRC RIS 2008-11, “NRC Regulatory Issue Summary 2008-11: Precautions to Protect Children Who May Come in Contact with Patients Released After Therapeutic Administration of Iodine-131”, Nuclear Regulatory Commission, May 2008.

⁵⁷ See Report Appendix, “Radiation Dose Calculations for I-131 Therapy Patients Released to a Hotel”

385 and easily understood oral and written instructions, and judges that the patient, or the patient's
386 parent or guardian, understands the instructions and is capable of complying with the recommended
387 precaution actions, then the dose to any other individual exposed to the I-131 therapy patient is
388 likely not to exceed 1 mSv even when released to a location other than a private residence.

389
390 The ICRP⁵⁸ suggested that a patient could “stay at a nonhospital living facility, such as a hotel,
391 for several days” when the patient's home situation would put the patient in close contact with
392 children due to physical or social constraints, because this “is less expensive than staying in a
393 hospital”. Initial research survey results conducted with voluntary respondents from the Thyroid
394 Cancer Survivors' Association indicated that most released patients in the U.S. go to a private
395 residence (approximately 94%) and only a few (approximately 5%) go to hotels⁵⁹. The
396 Subcommittee agrees that I-131 therapy patient release to a private residence should be encouraged,
397 and that licensees should carefully evaluate patient release to other locations and communicate to
398 the patient additional radiation safety precautions that may be appropriate for such locations.

399
400 The Subcommittee discussed management of dose to other individuals exposed to multiple
401 released patients as might occur with workers in a hotel near a major medical facility or workers in
402 a nursing home. The NRC's final rulemaking states that its medical experts “concluded that no
403 common nuclear medicine practice, be it diagnostic, therapeutic, or a combination of the two,
404 results in multiple large administrations that would be likely to cause the 5-millisievert (0.5-rem)
405 dose limit to be exceeded because of multiple administrations in a year”⁶⁰. The Subcommittee
406 extensively discussed patient release to hotels in regard to whether:

- 407
- 408 • dose management is adequate with current patient release instructions,
 - 409 • additional guidance and patient instructions are needed,
 - 410 • there should be added regulatory criteria, and
 - 411 • this dose management would be effectively accomplished by focusing only on I-131 therapy
412 patient release rather than trying to sum small doses from all radioactive material released
413 patients.
- 414

415 One Subcommittee member felt that no patients should be released to hotels or other similar
416 locations, and one Subcommittee member felt uneasy about allowing this release. Two
417 Subcommittee members felt that patients should be allowed to go a hotel, but that a licensee should,
418 by NRC guidance, track and control the number of released I-131 therapy patients planning to go to
419 specific hotels. Four Subcommittee members felt release to hotels was an acceptable option, and
420 there was no need to track or control release to specific hotels because the realistic projected dose to
421 others is small⁶¹. The different perspectives of the Subcommittee members on how best to assure
422 compliance with the applicable dose limits led us to conclude that the NRC should support a wider
423 discussion on this topic with the medical community and the public.

⁵⁸ ICRP Publication 94, “Release of Patients after Therapy with Unsealed Radionuclides”, International Commission on Radiological Protection, March 2004 – see paragraph (106), item (v).

⁵⁹ Vetter R, Van Nostrand D, Khorjekar G, et al, Presentation on “Use of a Patient Survey to Evaluate Compliance with and Quality of Instructions Given to Patients Treated with Radioiodine”, Annual Meeting of the Health Physics Society, Salt Lake City, Utah, June 27-July 1, 2010.

⁶⁰ 62 FR 4120: “Criteria for the Release of Individuals Administered Radioactive Material-Final Rule”, NRC Docket No. RIN 3150-AE41, January 29, 1997.

⁶¹ See Report Appendix, “Radiation Dose Calculations for I-131 Therapy Patients Released to a Hotel”

424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461

Annual Dose Limits versus Per-Release Dose Limits

The current 10 CFR 35.75 release criteria were developed in accordance with the NRC’s stated practice of implementing risk-informed performance-based regulations for licensees. The NRC appropriately recognized that licensees would only be able to judge “likely” doses to other individuals based on knowledge shared by patients of their post-release living circumstances and on the patients’ ability to follow instructions in maintaining these doses as low as reasonably achievable. Once the patient is released, the licensee no longer controls the patients’ actions, and patients are not accountable to NRC regulations. As stated in the final rulemaking for 10 CFR 35.75⁶²:

“The NRC is establishing a dose limit of 5 millisieverts (0.5 rem) total effective dose equivalent to an individual from exposure to the released patient for each patient release.”

The ICRP recommended dose constraint of a few mSv/episode “has often been inappropriately interpreted as a rigid annual dose limit”⁶³. The Subcommittee considered the consequences of changes to the current 10 CFR 35.75 release criteria, which apply to all diagnostic and therapeutic radioactive materials administered to patients and human research subjects, from a per-release limit to a rigid annual dose limit. The primary difficulty identified was the practicality of licensees tracking all doses to other individuals on an annual basis, potentially including those from multiple therapy administrations to the same patient in a single calendar year. The NRC concluded in their final rulemaking that the level of recordkeeping, even when limited to patient releases likely to exceed 0.1 mSv, was “an unnecessary burden”, and NRC clearly stated⁶²:

“Each patient release is to be treated as a separate event, and licensee knowledge of previous administrations is unnecessary.”

The NRC published a regulatory issue summary in 2008 which stated its intent to pursue rulemaking to change the 10 CFR 35.75 patient release criteria from dose limits to dose per year limits because the “presumption that patients receive single administrations of therapeutic doses in a given year, which is the basis used in developing the wording for the dose limit in 10 CFR 35.75, is no longer valid”⁶⁴. The RIS states NRC’s view of how licensees should manage patient release involving multiple administrations or applications in a single year. While the NRC explained that it would follow normal rulemaking procedures, including opportunity for public comment, this RIS created confusion as to whether the current 10 CFR 35.75 patient release criteria are per-release or annual dose limits⁶⁵.

⁶² 62 FR 4120: “Criteria for the Release of Individuals Administered Radioactive Material-Final Rule”, NRC Docket No. RIN 3150-AE41, January 29, 1997.

⁶³ ICRP Publication 94, “Release of Patients after Therapy with Unsealed Radionuclides”, International Commission on Radiological Protection, March 2004.

⁶⁴ NRC Regulatory Issue Summary 2008-07: “Dose Limit for Patient Release Under 10 CFR 35.75”, March 27, 2008.

⁶⁵ Prior to review of the 10 CFR 35.75 rulemaking notices in the Federal Register, polling of the Subcommittee members indicated that half of the members believed current release criteria were per-release dose limits and half believed the criteria were annual limits.

462 The Subcommittee reviewed and compared the Federal Register proposed rulemaking⁶⁶ and the
 463 final rulemaking⁶⁷ which established the current 10 CFR 35.75 patient release criteria. The NRC
 464 clearly stated in its proposed rulemaking that the patient release criteria would be annual dose
 465 limits. However, in the final rulemaking, the NRC changed the patient release criteria by dropping
 466 the annual limits and instead making the limits apply to each patient release. In regard to this
 467 change, the NRC stated⁶⁷,

468
 469 “Upon reconsideration, based on public comments and consultation with the ACMUI, an
 470 NRC medical consultant, and the NRC Visiting Medical Fellow, the NRC has decided to
 471 delete this requirement. A review of medical treatment practices revealed no common
 472 practice that would result in doses exceeding the 5 millisievert (0.5 rem) limit because of
 473 multiple administrations in the same year to the same patient. Without the need to account
 474 for the dose from multiple administrations, maintaining records for the many tens of
 475 thousands of patients released when their dose to an individual is likely to exceed 1
 476 millisievert (0.1 millisievert) becomes an unnecessary burden. The requirement to retain
 477 these records has therefore been deleted. Each patient release is to be treated as a separate
 478 event, and licensee knowledge of previous administrations is unnecessary.”
 479

480 There has been significant growth in the use of radioactive material medical procedures in the past
 481 20 years⁶⁸, and a few medical procedures, including a few I-131 therapy procedures, are
 482 administered to patients more than one time within a calendar year. However, exposure from
 483 multiple patients undergoing diagnostic procedures continues to be low in doses to other
 484 individuals. Exposure to a patient undergoing multiple I-131 therapies (2 to 3) in one year is likely
 485 to be a low dose to other individuals because of the patient following simple instructions for their
 486 release. Moreover if one applies the theory of linear no threshold radiation risk, there would be no
 487 difference in theoretical risk of radiation dose from exposure to an I-131 therapy patient receiving
 488 two therapies in one calendar year versus exposure to an I-131 therapy patient receiving a therapy
 489 per year in two calendar years.
 490

491 Based on the NRC conclusions documented in its final rulemaking⁶⁷ and lack of further
 492 rulemaking changes to the current 10 CFR 35.75 patient release criteria, the Subcommittee
 493 recommends the current patient release criteria should continue to be considered as per-release dose
 494 limits until modified by future rulemaking. Seven Subcommittee members believe that a new
 495 requirement for annualized dose limits could severely limit patients’ access to appropriate medical
 496 care at reasonable costs⁶⁹. These Subcommittee members conclude that the most effective and
 497 practical way to control the dose to other individuals from the release of patients administered
 498 radioactive materials is to support development of new guidance and other tools to assist: (a)

⁶⁶ 59 FR 30724: “Criteria for the Release of Patient Administered Radioactive Material, Proposed Rule”, NRC Docket No. RIN 3150-AE41, June 15, 1994.

⁶⁷ 62 FR 4120: “Criteria for the Release of Individuals Administered Radioactive Material-Final Rule”, NRC Docket No. RIN 3150-AE41, January 29, 1997.

⁶⁸ NCRP Report 160, “Ionizing Radiation Exposure of the Population of the United States”, National Council on Radiation Protection and Measurements, March 2009.

⁶⁹ One Subcommittee member believed that a dose limit would not be a true limit without an associated time frame. The remaining Subcommittee members believed strict adherence to an annual dose limit would severely limit access to medical care, and that the type and typical number of radioactive material medical procedures for a given patient do not result in excessive dose to other individuals in a calendar year.

499 licensees in assessing, carrying out, and documenting patient release; and (b) patients in
500 understanding and taking appropriate precautions for their specific living circumstances.

501
502 One Subcommittee member felt that the inconsistency and confusion over the per-release and
503 annual limit was due to the regulatory nature of the regulation. A per-event limit without an annual
504 limit allows an individual to receive multiple exposures. Although highly unlikely, this situation
505 would be allowable. Furthermore, an annual limit that is the same as a per-release limit is
506 duplicative, since the per-release limit would then be unnecessary. This one Subcommittee member
507 believes the simple solution would be to increase the annual limit for a caregiver who is exposed
508 more than once in a calendar year.

509 Petition to Return to Pre-1997 Release Criteria

510
511
512 The NRC was petitioned⁷⁰ to replace the current dose-based release criteria and to re-instate the
513 1986 10 CFR 35.75 release criteria⁷¹, widely known as the “30-mCi” rule. The NRC has also
514 received other requests to return to this old rule⁷². The Subcommittee finds no scientific merit in
515 returning to such activity-based release criteria, which have no identifiable scientific basis⁷³. The
516 Subcommittee maintains that dose-based release criteria are more scientifically rigorous than
517 activity-based criteria and better protect the public by basing patient releasability on the quantity,
518 dose, *directly* related to potential radiation hazard rather than on a quantity, activity, *indirectly*
519 related to this potential hazard. In the case of I-131 treatment of thyroid cancer, for example, the
520 administered I-131 is rapidly excreted (assuming a whole-body biological half-time of only about 2
521 days or less). In treating hyperthyroidism, however, 25 to 50% or more of the radioiodine localizes
522 in the thyroid, and that activity is cleared from the gland (and, in turn, the body) much more slowly,
523 with half-times of about 20 days or longer. Accordingly, the retained activity from the much higher
524 activity (typically greater than 100 mCi) administered to the thyroid cancer patient is rapidly
525 reduced to a lower activity than that retained by hyperthyroid patients (who typically receive about
526 10 mCi)⁷⁴. Thus, higher dose-rate irradiation of individuals persists longer for lower-activity
527 treatment of hyperthyroidism than for higher-activity treatment of thyroid cancer, illustrating the
528 fallacy of an idea that activity-based release criteria (i.e. the “30-mCi” rule) is more protective of
529 public safety^{74,75,76}.

530
531 In fact, the 30-mCi rule is a special case of the 1997 release criteria, based on I-131 with the
532 following conditions:

533

⁷⁰ 70 FR 75752, “Peter G. Crane; Receipt for Rulemaking”, NRC Docket No. PRM-35-18, December 21, 2005.

⁷¹ 51 FR 36932, “Medical Use of Radioactive Material-Final Rule”, Nuclear Regulatory Commission, October 16, 1986.

⁷² “Radioactive Roulette: How the Nuclear Regulatory Commission’s Cancer Patient Radiation Rules Gamble with Public Health and Safety”, A report by the Staff of Edward J. Markey (D-MA), Chairman, Subcommittee on Energy and Environment, Energy and Commerce Committee, U.S. House of Representatives, March 18, 2010.

⁷³ Siegel JA, “Tracking the Origin of the NRC 30-mCi Rule”, J Nucl Med. 2000;41:10-16N.

⁷⁴ ICRP Publication 94, “Release of Patients after Therapy with Unsealed Radionuclides”, International Commission on Radiological Protection, March 2004.

⁷⁵ IAEA Safety Reports Series No. 63, “Release of Patients after Radionuclide Therapy”, International Atomic Energy Agency, 2009.

⁷⁶ See Report Appendix, “Radiation Dose Calculations for I-131 Therapy Patients Released to a Hotel”.

- 534 • Using the physical half-life instead of the effective half-life, ignoring biological
535 elimination of the radionuclide;
536 • Ignoring the attenuation of the radiation by the patient;
537 • Using the default occupancy of 0.25 rather than a value based on actual patient behavior
538 information.
539

540 The 30-mCi rule also represented a “per-release” limit. Returning to the old rule simply would
541 ignore physical principles as well as consideration of actual patient behavior in different living
542 circumstances. Change from the 30-mCi rule to the current 10 CFR 35.75 patient release criteria in
543 no way weakened the NRC rules.
544

545 NRC policy was not intended to intrude on the practice of medicine⁷⁷, yet evidence exists that
546 prior to adopting the 1997 risk based release criteria, the former activity-based release criteria
547 adversely impacted the practice of medicine and patient care by limiting patients to only 30-mCi
548 administered activities simply to allow immediate patient release. This practice essentially
549 fractionates the patient’s therapy dose and reduced the effectiveness of therapy. In some countries
550 where activity-based release criteria are still used, patients are effectively denied therapy for as long
551 as one year because of lack of hospital rooms for overnight accommodation. The Subcommittee
552 commends the NRC for adopting the current-risk-based criteria.
553

554 Developing Updated Guidance in Support of Patient Release Dose Controls

555 The NRC guidance to licensees on patient release criteria⁷⁸ was based on dose calculation
556 methods and assumptions that are overly conservative and outdated. The Subcommittee
557 recommends that the NRC, with assistance from experts, update the patient release guidance using
558 reasonable assumptions based on an expanded list of radionuclides used in medicine, current
559 radiopharmaceutical biokinetics information, and reported dose measurements from patients.
560 Computer-based modes of communications, data gathering, and data processing should be used to
561 develop tools and accrue data for guidance of licensees in:
562

- 563 • assessing various living situations, including patient release to other locations (such as
564 hotels, public transport, public events),
565 • calculating realistic radiation dose to others,
566 • choosing realistic precautions for patients to take,
567 • instructing patients on these precautions and specific applications, and
568 • documenting compliance with the patient release criteria.
569

570 During this review, the Subcommittee found many scholarly efforts which have advanced
571 understanding and communication of real-world situations that impact patient release decisions and
572 perceptions. The NRC should support research activities to better identify what aspects of patient
573 release have realistic impact on doses to other individuals. As examples, the following efforts
574 provide insights into various aspects of patient release.
575

⁷⁷ 65 FR 47654, “Medical Use of Byproduct Material; Policy Statement, Revision”, Nuclear Regulatory Commission, August 3, 2000.

⁷⁸ NRC Regulatory Guide 8.39, “Release of Patients Administered Radioactive Materials”, Nuclear Regulatory Commission, April 1997.

- 576
- 577 • Measurements of radiation exposure to household members from released patients⁷⁹
- 578 • Surveys of patients and caregivers to determine understanding of and adherence to patient
- 579 release instructions⁸⁰
- 580 • Communication tools to help convey personalized instructions to patients⁸¹
- 581 • Credible websites providing objective, scientific information about radiation⁸²
- 582 • Medical protocol enhancements for patient release⁸³
- 583

584 Patients want access to the best health care. And while release of the I-131 therapy patient is

585 most often the focus of evaluating the potential hazard to others, the I-131 patient should not be

586 treated unfairly by virtue of need for I-131 therapy. Well-informed patients are self-motivated and

587 sensitive to the fact that they are radioactive for a period of time, excreting radioactivity, and will

588 typically do as much as possible to reduce potential exposures to family, caregivers, and other

589 members of the general public. They need to be reassured that their medical procedure with

590 radioactive material is safe for themselves, their family members and their caregivers, and that they

591 do not represent a source of harmful radiation exposure to members of the public. Any new NRC

592 guidance should be developed with the assistance of experts involved with patient release⁸⁴, and

593 focus on improved patient counseling rather than excessive controlling or monitoring of the patient.

594

595

596 **Subcommittee Conclusions on Patient/Human Research Subject Release Issues**

597

598 The Subcommittee commends the NRC for its leadership role in developing and implementing

599 practical regulatory control of the use of radioactive materials in patients which appropriately

600 applies the three fundamental radiation protection principles of justification, optimization and

601 limits. Benefits from medical use of radioactive materials are many and well-recognized,

602 improving the health and lives of millions of people in the U.S. These benefits far exceed the small

603 theoretical risks associated with exposure from released patients.

604

⁷⁹ Grigsby PW, Siegel BA, Baker S, & Eichling, JO. "Radiation exposure from outpatient radioactive iodine (I-131) therapy for Thyroid Carcinoma". JAMA. 2000;283:2272-2274.

⁸⁰ Vetter R, Van Nostrand D, Khorjekar G, et al, Presentation on "Use of a Patient Survey to Evaluate Compliance with and Quality of Instructions Given to Patients Treated with Radioiodine", Annual Meeting of the Health Physics Society, Salt Lake City, Utah, June 27-July 1, 2010.

⁸¹ Freidman MI, Ghesani M, "Interactive Software Automates Personalized Radiation Safety Plans for Na¹³¹I Therapy", Health Physics (83 Supplement 5:S71-S84), November 2002.

⁸² "Radiation Answers: Answers to Questions About Radiation and You", www.radiationanswers.org, supported by the Health Physics Society.

⁸³ Khorjekar G, Van Nostrand D, Vetter R, et al, Poster on "The Relationship of Several Factors and Vomiting After Outpatient I-131 Therapy in Patients with Well-Differentiated Thyroid Cancer", Society of Nuclear Medicine Annual Meeting, Salt Lake City, Utah, June 5-9, 2010.

⁸⁴ The Subcommittee members differed in their opinions on methods needed to best counsel multiple patients in managing release to the same location, but agreed that it is essential for the NRC to work with the medical community and the public to develop reasonable and effectual guidance which minimizes impacts on patient access to these medical procedures.

605 The Health Physics Society⁸⁵ recently updated their position statement regarding radiation
606 risk⁸⁶, and stated:

607
608 “In accordance with current knowledge of radiation health risks, the Health Physics Society
609 recommends against quantitative estimation of health risks below an individual dose of 5
610 rem in one year or a lifetime dose of 10 rem above that received from natural sources.
611 Doses from natural background radiation in the United States average about 0.3 rem per
612 year. A dose of 5 rem will be accumulated in the first 17 years of life and about 25 rem in a
613 lifetime of 80 years. Estimation of health risk associated with radiation doses that are of
614 similar magnitude as those received from natural sources should be strictly qualitative and
615 encompass a range of hypothetical health outcomes, including the possibility of no adverse
616 health effects at such low levels.

617
618 There is substantial and convincing scientific evidence for health risks following high-dose
619 exposures. However, below 5–10 rem (which includes occupational and environmental
620 exposures), risks of health effects are either too small to be observed or are nonexistent.”

621
622 Ongoing research efforts are exploring the effects of low-dose radiation exposures^{87,88} and
623 examining whether health impacts exist in populations exposed to low levels of radiation^{89,90,91,92}.

624
625 Regulatory decision-making is ultimately a politically based national policy discussion⁹³ which
626 is shaped by opinions sometimes based on the perception rather than the reality of risk⁹⁴. The NRC
627 remains an important leader in this national discourse⁹⁵. In light of limited health care resources, it
628 is increasingly important that regulations serve not only to protect society from *real* hazards, but

⁸⁵ The Health Physics Society is a nonprofit scientific professional organization whose mission is excellence in the science and practice of radiation safety. Since its formation in 1956, the Society has grown to approximately 6,000 scientists, physicians, engineers, lawyers, and other professionals representing academia, industry, government, national laboratories, the Department of Defense, and other organizations. Society activities include encouraging research in radiation science, developing standards, and disseminating radiation safety information. Society members are involved in understanding, evaluating, and controlling the potential risks from radiation relative to the benefits.

⁸⁶ HPS PS010-2, “Radiation Risk in Perspective”, Position Statement of the Health Physics Society, revised July 2010.

⁸⁷ Brooks AL, “Developing a Scientific Basis for Radiation Risk Estimates: Goal of the DOE Low Research Program”, Health Physics (85:85-93), July 2003.

⁸⁸ Averbeck D, “Does Scientific Evidence Support a Change from the LNT Model for Low-Dose Radiation Risk Extrapolation?”, Health Physics (97:493-504), November 2009.

⁸⁹ Shore RE, “Low-Dose Radiation Epidemiology Studies: Status and Issues”, Health Physics (97:481-486), November 2009.

⁹⁰ Dickman PW, et. al., “Thyroid Cancer Risk After Thyroid Examination with I-131: a Population-Based Cohort Study in Sweden”, Int. J. Cancer: 106, 580-587 (2003).

⁹¹ Ghiassi-nejad M, et al, “Very High Background Radiation Areas of Ramsar, Iran: Preliminary Biological Studies”, Health Physics (82:87-93), January 2002.

⁹² Nair RRR, et al, “Background Radiation and Cancer Incidence in Kerala, India-Karunagappally Cohort Study”, Health Physics (96:55-66), January 2009.

⁹³ Locke P, “Incorporating Information from the U.S. Department of Energy Low-Dose Program into Regulatory Decision-Making: Three Policy Integration Challenges”, Health Physics (97:510-515), November 2009.

⁹⁴ Jenkins-Smith HC, Silva CL, Murray C, “Beliefs about Radiation: Scientists, the Public and Public Policy”, Health Physics (97:519-527), November 2009.

⁹⁵ Tenforde TS, Brooks AL, “Perspectives of U.S. Government Agencies on the Potential Role of Greater Scientific Understanding of Low-Dose Radiation Effects in Establishing Regulatory Health Protection Guidance”, Health Physics (97:516-518), November 2009.

629 that they also be based on realistic projections of the severity and likelihood, and on consideration
630 of the actual costs, financial and otherwise, from overly cautious and potentially intrusive
631 regulations. For radionuclide therapy that has been shown to be a safe, effective, and financially
632 viable treatment for certain cancers and other serious diseases, patient release criteria and relevant
633 regulations based on realistic dose projections are both conducive to public safety and promote
634 access to and affordability of such therapy. The Subcommittee affirms that the current dose-based
635 release criteria 10 CFR 35.75 meet these essential benchmarks.
636

637 The Subcommittee therefore concludes that the current 10 CFR 35.75 release criteria
638 appropriately balance public safety with patient access to efficacious and cost-effective medical
639 treatment. The Subcommittee recommends that the NRC gather scientific data on patient behavior
640 and understanding of instructions to determine the most effective instructions to enhance licensee
641 communication and documentation of patient release, and to promote patient understanding. The
642 Subcommittee further recommends that the NRC update patient release guidance, with assistance
643 from experts, to include current information on actual radiopharmaceutical biokinetics and
644 calculated or measured patient dose rates, and provide guidance for release scenarios to other
645 locations other than private residences (such as hotels, public transport, public events).

646 **Appendix⁹⁶ – Radiation Dose Calculations for I-131 Therapy Patients Released to a Hotel**
 647
 648

649 The Subcommittee conducted a scientific analysis of radiation doses that might be received
 650 by hotel workers in the event that an iodine-131 (I-131) therapy patient, appropriately released
 651 from a medical institution, chose to stay in a hotel immediately following the release. We show
 652 for four scenarios what the radiation doses to hotel workers and other guests could be under
 653 different sets of parameters. The four scenarios are labeled *unrealistic* (representing an
 654 improbable, worst-case scenario), *highly unlikely* (representing a doubtful scenario, rarely
 655 occurring), *conservative* (representing a possible scenario, not likely to occur), and *realistic*
 656 (representing a more likely scenario for a typical patient). The four scenarios involve release to a
 657 hotel of (1) an I-131 cancer therapy patient (Table 1), and (2) an I-131 hyperthyroid therapy
 658 patient (Table 2). The assumptions and parameters used for each scenario are described in each
 659 table.

660
 661 Published scientific literature indicates that radiation doses to non-patients from iodine-131
 662 patients released after therapy may consist of two components: (1) external radiation exposure
 663 received by standing in close proximity to the patient, and (2) the intake of I-131 contamination
 664 from I-131 that leaves the patient in excreta or sweat. The literature shows that an individual's
 665 radiation dose from the uptake of I-131 contamination is far less significant (less than 10%) than
 666 the radiation dose received from external exposure to the patient^{97,98,99,100}. Radiation
 667 measurements have shown that internal contamination of family members from radioactive
 668 patients may only be something on the order of one-millionth of the activity administered to the
 669 patient. Therefore, the potential radiation dose to a family member or hotel worker from
 670 internalized contamination left by a released I-131 patient can only be far below that which is
 671 possible from external doses^{101,102,103,104,105,106} (also see Table 3). In addition, the likelihood of

⁹⁶ Appendix to the Advisory Committee on the Medical Use of Isotopes (ACMUI) Patient Release Subcommittee Report, December 6, 2010 draft.

⁹⁷ NCRP Commentary No. 11, "Dose Limits for Individuals Who Receive Exposure from Radionuclide Therapy Patients", National Council on Radiation Protection and Measurements, February 1995.

⁹⁸ NRC NUREG-1492, "Regulatory Analysis on Criteria for the Release of Patients Administered Radioactive Materials, Final Report", by Stewart Schneider and Stephen A. McGuire, Nuclear Regulatory Commission, April 1996.

⁹⁹ ICRP Publication 94, "Release of Patients after Therapy with Unsealed Radionuclides", International Commission on Radiological Protection, March 2004.

¹⁰⁰ IAEA Safety Reports Series No. 63, "Release of Patients after Radionuclide Therapy", International Atomic Energy Agency, 2009.

¹⁰¹ Buchan RCT, Brindle JM. "Radioiodine therapy to outpatients—the contamination hazard". *Br J Radiol* 43:479–482; 1970.

¹⁰² Hammond N, Jacobson A. An effective method to reduce the exposure to families of radioiodine therapy patients. *Health Phys.* 1982;43:89-172.

¹⁰³ Jacobson AP, Plato PA, Toeroek D. "Contamination of the home environment by patients treated with iodine-131: initial results". *Am J Public Health* 68:230–235; 1978.

¹⁰⁴ Plato P, Jacobson A, Homann S. "In vivo thyroid monitoring for iodine-131 in the environment". *Inter J Applied Radiat Isotopes.* 1976;27:539-545.

¹⁰⁵ Toeroek D, Jacobson A, Plato P. "Radiation protection of families of radioactive patients". *Health Phys.* 1978;35:911-912.

¹⁰⁶ Chandra R, Marshall C. "Radioiodine therapy to out-patients - The contamination hazard (Letter)". *Br J Radiol.* 1971;44:557.

672 an intake following intermittent exposure to I-131 contamination of toilets and bedding is very
673 small for both immediate family members and for hotel guests or workers¹⁰⁷. Even if a hotel
674 worker were *not* to wear gloves while cleaning a released I-131 therapy patient's room, the risk
675 of internalization of I-131 radioactivity remains low – as indicated by the data in the references
676 in Footnote 101 and Table 3.

677
678 Despite the use of these overly-cautious assumptions and parameters used in Tables 1 and 2,
679 the highest projected dose to a hotel housekeeper from a released cancer therapy patient is less
680 than 100 mrem. For the case of a released hyperthyroid patient treated for immediate release
681 under the 30-mCi rule, where the amount of I-131 administered is 17% of the amount
682 administered to the cancer therapy patient (Table 2), the three-day projected doses to a hotel
683 housekeeper are 67% of that from the released cancer therapy patient.

684
685 The realistic projected doses to hotel workers are very low. To give a perspective of how
686 safe these projected doses are, the average U.S. dose from natural background radiation is 310
687 mrem per year¹⁰⁸, or 0.85 mrem per day. The highest realistic hotel worker dose of 1.2 mrem
688 would be equivalent to an extra 1.4 days of natural background radiation. The highest realistic
689 guest dose of 22 mrem would be equivalent to an extra 26 days of natural background radiation.

690
691 Use of patient-specific parameters in conjunction with realistic assumptions of behavior by
692 the patient, hotel workers and other guests should be used when calculating a particular patient
693 release.

¹⁰⁷ Personal correspondence from M.G. Stabin, Ph.D., CHP.

¹⁰⁸ NCRP Report No. 160, "Ionizing Radiation Exposure of the Population of the United States", National Council on Radiation Protection and Measurements, March 2009.

TABLE 1 –Radiation Dose Calculations to Hotel Workers and Guests from an I-131 Cancer Therapy Patient

- 175 mCi ¹³¹I-iodide administered to a post-thyroidectomy thyroid cancer patient
- Doses calculated assuming point source*: patient self-shielding** (0.13 mrem-m²/h-mCi); laundry no shielding (0.22 mrem-m²/h-mCi)
- Total-body effective time-activity function*: $0.95 e^{(0.693/0.32 \text{ day}) t} + 0.05 e^{(-0.693/7.3 \text{ day}) t}$
- Mean distance from patient to guest in adjoining room is 2.2 m (based on mid-point of 80 inch long beds + 6 inch wall), assuming no shielding provided by walls between rooms, and assuming head to head exposure equals mid-body to mid-body exposure
- Dose contribution of possible internal radioactive contamination is considered minor and not included

| Assumptions and Parameters*** | Unrealistic | | | Highly Unlikely | | | Conservative | | | Realistic | | |
|---|--|----|----|--------------------|----|----|--------------------|-----|-----|--|-------|------|
| | Time (in days) Patient Remained in Hotel | | | | | | | | | | | |
| Cohort | 1 | 2 | 3 | 1 | 2 | 3 | 1 | 2 | 3 | 1 | 2 | 3 |
| | Radiation Dose to Cohort (in mrem) | | | | | | | | | | | |
| Hotel Housekeeper | 69 | 83 | 91 | 35 | 43 | 47 | 14 | 17 | 18 | 0.90 | 1.1 | 1.2 |
| Hotel Laundry Worker | 39 | 47 | 52 | 16 | 19 | 21 | 3.9 | 4.7 | 5.2 | 0.078 | 0.095 | 0.10 |
| Non-Housekeeping/Non-Laundry Hotel Worker or Hotel Guest in Non-Adjoining Room | 30 | 36 | 39 | 20 | 24 | 26 | 10 | 12 | 13 | 0.83 | 0.99 | 1.1 |
| Hotel Guest in Room Adjoining that of Patient | 54 | 65 | 71 | 40 | 48 | 53 | 26 | 32 | 34 | 17 | 21 | 22 |
| Parameters | | | | | | | | | | | | |
| Remaining activity in patient excreted into bed linens at midpoint of each day | 50% per day | | | 20% per day | | | 5% per day | | | 0.1% per day (bath linens & cleaning only) | | |
| Time hotel housekeeper and laundry worker each hold contaminated linens (0.3 m away) | 30 minutes per day | | | 20 minutes per day | | | 10 minutes per day | | | 10 minutes per day | | |
| Time hotel housekeeper, other workers (except laundry), and other guests are 1 meter from patient | 3 hours per day | | | 2 hours per day | | | 1 hour per day | | | 5 minutes per day | | |
| Additional time patient and other hotel guest in adjoining room are both in their respective beds | 12 hours per day | | | 10 hours per day | | | 8 hours per day | | | 8 hours per day | | |

* Values used are from NRC Regulatory Guide 8.39

** Patient self-shielding value from SPARKS, R.B., SIEGEL, J.A. and WAHL, R.L. (1998). "The need for better methods to determine release criteria for patients administered radioactive material," Health Phys. **75**(4), 385–388.

***These assumptions and parameters should be adjusted for patient-specific situations, considering patient release instructions, to calculate realistic doses.

TABLE 2 – Radiation Dose Calculations to Hotel Workers and Guests from an I-131 Hyperthyroid Patient

- 29.9 mCi ¹³¹I-iodide administered to a hyperthyroid patient
- Doses calculated assuming point source*: patient self-shielding** (0.13 mrem-m²/h-mCi); laundry no shielding (0.22 mrem-m²/h-mCi)
- Total-body effective time-activity function*: $0.20 e^{(0.693/0.32 \text{ day}) t} + 0.80 e^{(-0.693/5.2 \text{ day}) t}$
- Mean distance from patient to guest in adjoining room is 2.2 m (based on mid-point of 80 inch long beds + 6 inch wall), assuming no shielding provided by walls between rooms, and assuming head to head exposure equals mid-body to mid-body exposure
- Dose contribution of possible internal radioactive contamination is considered minor and not included

| Assumptions and Parameters*** | Unrealistic | | | Highly Unlikely | | | Conservative | | | Realistic | | |
|---|--|----|----|--------------------|----|----|--------------------|-----|-----|--|-------|-------|
| | 1 | 2 | 3 | 1 | 2 | 3 | 1 | 2 | 3 | 1 | 2 | 3 |
| | Time (in days) Patient Remained in Hotel | | | | | | | | | | | |
| | Radiation Dose to Cohort (in mrem) | | | | | | | | | | | |
| Cohort | | | | | | | | | | | | |
| Hotel Housekeeper | 25 | 44 | 61 | 12 | 22 | 31 | 4.7 | 8.5 | 12 | 0.30 | 0.54 | 0.74 |
| Hotel Laundry Worker | 15 | 27 | 37 | 5.9 | 11 | 15 | 1.5 | 2.7 | 3.7 | 0.029 | 0.053 | 0.074 |
| Non-Housekeeping/Non-Laundry Hotel Worker or Hotel Guest in Non-Adjoining Room | 10 | 17 | 24 | 6.4 | 12 | 16 | 3.2 | 5.8 | 8.0 | 0.27 | 0.48 | 0.67 |
| Hotel Guest in Room Adjoining that of Patient | 18 | 32 | 44 | 13 | 24 | 33 | 8.5 | 15 | 21 | 5.6 | 10 | 14 |
| Parameters | | | | | | | | | | | | |
| Remaining activity in patient excreted into bed linens at midpoint of each day | 50% per day | | | 20% per day | | | 5% per day | | | 0.1% per day (bath linens & cleaning only) | | |
| Time hotel housekeeper and laundry worker each hold contaminated linens (0.3 m away) | 30 minutes per day | | | 20 minutes per day | | | 10 minutes per day | | | 10 minutes per day | | |
| Time hotel housekeeper, other workers (except laundry), and other guests are 1 meter from patient | 3 hours per day | | | 2 hours per day | | | 1 hour per day | | | 5 minutes per day | | |
| Additional time patient and other hotel guest in adjoining room are both in their respective beds | 12 hours per day | | | 10 hours per day | | | 8 hours per day | | | 8 hours per day | | |

* Values used are from NRC Regulatory Guide 8.39

** Patient self-shielding value from SPARKS, R.B., SIEGEL, J.A. and WAHL, R.L. (1998). "The need for better methods to determine release criteria for patients administered radioactive material," Health Phys. **75**(4), 385–388.

***These assumptions and parameters should be adjusted for patient-specific situations, considering patient release instructions, to calculate realistic doses.

Table 3 – Summary Table of Family Doses from Buchan Reference in Footnote 101

Summary of Radiation Dosimetry for Family Members of I131-Treated Thyroid Patients

| Disease Treated | Exposed Cohort | Maximum Recommended Annual Effective Dose | | Radiation Precautions | Equivalent Dose per Unit Patient Total Body Activity at Discharge | | | | | | Effective Dose per Unit Activity at Discharge | | | | | | Patient Activity at Discharge for Maximum Recommended Effective Dose | | | | | |
|-----------------|------------------|---|------|-----------------------------|---|----------|--------------------|---------------|--------------------|----------|---|----------|--------------------|----------|--------------------|----------|--|------------|---------|--------|----|----|
| | | μSv | mrem | | Total Body | | Thyroid | | Total | | Total Body (non-Thyroid) | | Thyroid | | Total | | | | | | | |
| | | | | | $\mu\text{Sv/MBq}$ | mrem/mCi | $\mu\text{Sv/MBq}$ | mrem/mCi | $\mu\text{Sv/MBq}$ | mrem/mCi | $\mu\text{Sv/MBq}$ | mrem/mCi | $\mu\text{Sv/MBq}$ | mrem/mCi | $\mu\text{Sv/MBq}$ | mrem/mCi | $\mu\text{Sv/MBq}$ | mrem/mCi | MBq | mCi | | |
| Hyperthyroidism | Children | 1,000 | 100 | None / Minimal ^a | Minimum | 0.398 | 1.47 | <i>0.0789</i> | <i>0.292</i> | 0.477 | 1.76 | 0.378 | 1.40 | 0.0238 | 0.0881 | 0.402 | 1.48 | Best case | 2,490 | 67.4 | | |
| | | | | | Maximum | 0.920 | 3.42 | <i>35.1</i> | <i>204</i> | 56.0 | 207 | 0.874 | 3.25 | 2.80 | 10.4 | 3.68 | 13.6 | Worst case | 272 | 7.34 | | |
| | | | | | Median | 0.733 | 2.71 | <i>0.47</i> | <i>1.74</i> | 1.20 | 4.45 | 0.696 | 2.57 | 0.0602 | 0.223 | 0.757 | 2.80 | Median | 1,320 | 35.8 | | |
| | | | | | Minimum | 0.0611 | 0.300 | 0.0789 | 0.292 | 0.160 | 0.592 | 0.0770 | 0.285 | 0.00800 | 0.0296 | 0.0650 | 0.315 | Best case | 11,800 | 318 | | |
| | | | | | Maximum | 10.5 | 38.8 | 35.1 | 204 | 65.6 | 243 | 9.98 | 36.9 | 3.28 | 12.1 | 13.3 | 49.0 | Worst case | 76 | 2.04 | | |
| | | | | | Median | 1.99 | 7.35 | 0.47 | 1.74 | 2.46 | 9.09 | 1.89 | 6.98 | 0.123 | 0.455 | 2.01 | 7.44 | Median | 498 | 13.4 | | |
| | Adults / Spouses | 5,000 | 500 | None / Minimal | Minimum | 0.241 | 0.89 | 3 | 1.11 | 3.24 | 2.00 | 0.229 | 0.85 | 0.162 | 0.100 | 0.391 | 0.946 | Best case | 19,600 | 529 | | |
| | | | | | Maximum | 22.2 | 82.2 | 11.7 | 43.3 | 33.9 | 126 | 21.09 | 78.09 | 1.70 | 6.28 | 22.8 | 84.4 | Worst case | 219 | 5.93 | | |
| | | | | | Median | 4.83 | 17.9 | 1.88 | 7.00 | 6.71 | 24.9 | 4.59 | 17.01 | 0.336 | 1.25 | 4.92 | 18.3 | Median | 1,010 | 27.4 | | |
| | | | | | Minimum | 1.43 | 5.28 | 0.478 | 1.77 | 1.91 | 7.05 | 1.36 | 5.016 | 0.0954 | 0.353 | 1.45 | 5.37 | Best case | 3,450 | 93.1 | | |
| | | | | | Maximum | 13.1 | 48.3 | 10.2 | 37.6 | 23.3 | 85.9 | 12.44500 | 45.9 | 1.17 | 4.30 | 13.6 | 50.2 | Worst case | 369 | 10.0 | | |
| | | | | | Median | 1.79 | 6.64 | 2.03 | 7.52 | 3.82 | 14.2 | 1.70 | 6.31 | 0.191 | 0.708 | 1.89 | 7.02 | Median | 2,640 | 71.3 | | |
| Thyroid Cancer | Children | 1,000 | 100 | None / Minimal ^b | Minimum | 0 | 0.00 | <i>0.135</i> | <i>0.500</i> | 0.135 | 0.500 | 0.00 | 0.00 | 0.00675 | 0.0250 | 0.00675 | 0.0250 | Best case | 148,000 | 4,000 | | |
| | | | | | Maximum | 1.38 | 5.36 | <i>0.252</i> | <i>0.933</i> | 1.63 | 6.29 | 1.31 | 5.09 | 0.0816 | 0.315 | 1.39 | 5.41 | Worst case | 684 | 18.5 | | |
| | | | | | Median | 0.816 | 3.02 | <i>0.194</i> | <i>0.717</i> | 1.01 | 3.74 | 0.775 | 2.87 | 0.0505 | 0.187 | 0.826 | 3.06 | Median | 1,210 | 33 | | |
| | | | | | Minimum | NA | NA | 0.135 | 0.500 | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA |
| | | | | | Maximum | 0.0274 | 0.100 | 0.252 | 0.933 | 0.279 | 1.03 | 0.0260 | 0.0950 | 0.0140 | 0.0517 | 0.0400 | 0.147 | Worst case | 126,000 | 3,410 | | |
| | | | | | Median | NA | NA | 0.194 | 0.717 | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | |
| | Adults / Spouses | 5,000 | 500 | None / Minimal ^c | Minimum | 0.0137 | 0.0507 | <i>0.108</i> | <i>0.400</i> | 0.122 | 0.451 | 0.0130 | 0.0482 | 0.00609 | 0.0225 | 0.0191 | 0.0707 | Best case | 262,000 | 7,070 | | |
| | | | | | Maximum | 0.258 | 0.955 | <i>0.108</i> | <i>0.400</i> | 0.366 | 1.36 | 0.245 | 0.907 | 0.0183 | 0.0678 | 0.263 | 0.975 | Worst case | 19,000 | 513 | | |
| | | | | | Median | 0.0655 | 0.316 | <i>0.108</i> | <i>0.400</i> | 0.194 | 0.716 | 0.0812 | 0.300 | 0.00968 | 0.0358 | 0.0909 | 0.336 | Median | 55,100 | 1,490 | | |
| | | | | | Minimum | 0 | 0 | <i>0.108</i> | <i>0.400</i> | 0.108 | 0.400 | 0.00 | 0.00 | 0.00540 | 0.0200 | 0.00540 | 0.0200 | Best case | 925,000 | 25,000 | | |
| | | | | | Maximum | 2.27 | 8.400 | <i>0.108</i> | <i>0.400</i> | 2.38 | 8.80 | 2.16 | 7.98 | 0.119 | 0.440 | 2.28 | 8.42 | Worst case | 2,220 | 59.4 | | |
| | | | | | Median | 0.46 | 1.700 | <i>0.108</i> | <i>0.400</i> | 0.568 | 2.10 | 0.437 | 1.62 | 0.0284 | 0.105 | 0.465 | 1.72 | Median | 10,800 | 290.7 | | |

^a In the absence of available data for the thyroid-to-thyroid dose equivalent from internalized radioiodine in children of I131-treated hyperthyroid patients where radiation precautions were not observed, the corresponding data in children of I131-treated hyperthyroid patients where radiation precautions were observed were used (indicated by the italicized entries).

^b In the absence of available data for the thyroid-to-thyroid dose equivalent from internalized radioiodine in children of I131-treated thyroid cancer patients where radiation precautions were not observed, the corresponding data in children of I131-treated thyroid cancer patients where radiation precautions were observed were used (indicated by the italicized entries).

^c NA = Not Available

^d Data for the thyroid-to-thyroid dose equivalent from internalized radioiodine in adults / spouses of I131-treated thyroid cancer patients are available for only one subject, a subject for whom radiation precautions were observed. These data were therefore used as the median, minimum, and maximum and the minimum and maximum thyroid-to-thyroid dose equivalents values for adults / spouses of I131-treated thyroid cancer patients where radiation precautions were and were not observed, respectively (indicated by the italicized entries).